

LAB

Tel : 12345678

Address PUNE, INDIA

Patient Details

Name : DUMMY PATIENT

Date of Birth : 6 Sep 1973

Age : 50 years/Female

Contact Number : 1111111111 %>

Referred By :

Report Details

Name : Pregnancy Glucose Tolerance Test

Sampling Received At :

VERIFIED BY DR.XYZ ABC at 1 pm.

Name	Result	Normal Range
Pregnancy GTT - Plasma Glucose(Fasting) (mg/dL)	88	70-92
Pregnancy GTT - Plasma Glucose(1 hr) (mg/dL)	170	90-180
Pregnancy GTT - Plasma Glucose(2 hours) (mg/dL)	150	70-153

ALL PATHOFAST LAB REPORTS ARE SIGNED BY UNIQUE QR-CODES TO PREVENT FORGERY. SCAN THE QR-CODE NEXT TO THE SIGNATURE BELOW -> COMPARE THE PATIENT NAME DISPLAYED.

Disclaimer

This is a sample report, on a dummy patient, signed by a dummy doctor. The actual report format may vary and you are requested not to compare with the same.

- 'Sample Received At' refers to the date and time of the first sample collection done for a patient, for a given set of tests. In case of subsequent collections(eg in case of Fasting and PP, suppression or stimulation tests) the time may not reflect the actual time of collection.
- Methods of measurement are mentioned together at the bottom of each report, test units to the right of each test name.All reports are subject to clinical correlation and not intended for forensic purposes.

LAB

Tel : 12345678

Address PUNE, INDIA

Patient Details

Name : DUMMY PATIENT

Date of Birth : 6 Sep 1973

Age : 50 years/Female

Contact Number : 111111111 %>

Referred By :

Report Details

Name : Pregnancy Glucose Tolerance Test

Sampling Received At :

VERIFIED BY DR.XYZ ABC at 1 pm.

Authorized By



Dr. ABC XYZ
(DOCTOR)

Registration Number : 123456

Test Methods

1. GOD-POD

Notes, Interpretation & Interfering Factors

1. Fasting period of 8-14 hours is mandatory
2. International Association of Diabetes and Pregnancy Groups (IADPG) guidelines are used in our lab.
3. As per these guidelines, fasting, 1 hour and 2 hour samples are to be considered, after the administration of 75gm of glucose for the purpose of a gestational oral glucose tolerance test.
4. For the three time periods, Fasting , 1 hour and 2 hour : threshold values of 92 mg/dl, 180 mg/dl and 153 mg/dl, are used as cut-off values.
5. The IADPG recommends that all women should undergo a fasting PG and HBA1C during their first perinatal visit. A full GTT is recommended between 24-28 weeks of gestation with the above mentioned thresholds.
6. One or more value above the thresholds, is considered diagnostic of GDM(gestations diabetes mellitus)
7. There exist many different guidelines for interpretation and performance of the GTT, like ACOG, WHO, ADA , each with different normal ranges, glucose loading dose and sampling intervals. If your physician requires a specific guideline to be followed, please indicate the same at the time of the test.

Disclaimer

This is a sample report, on a dummy patient, signed by a dummy doctor. The actual report format may vary and you are requested not to compare with the same.

1. 'Sample Received At' refers to the date and time of the first sample collection done for a patient, for a given set of tests. In case of subsequent collections(eg in case of Fasting and PP, suppression or stimulation tests) the time may not reflect the actual time of collection.
2. Methods of measurement are mentioned together at the bottom of each report, test units to the right of each test name.All reports are subject to clinical correlation and not intended for forensic purposes.